

# **BLOOD CULTURE COLLECTION PROCEDURE KIT**

Including Sterile Initial Specimen Diversion Tube

## All together now.

sub dermally are separated and sequestered from augments accuracy, as bacteria that may be present 3-pocket design contains everything you need to optimize your efficiency with the new Blood Culture with C-Quest Blood Culture length and cost of hospital stays are minimized enhanced, while the use of antibiotics, as well as the results. Your proper diagnosis and treatment are methods that demonstrate higher false-positive test than previous options with non-diversion collection positive results. In addition, C-Quest Kits cost no more non-contaminated blood to virtually eliminate false collection. Plus, a sterile initial specimen diversion tube perform a safe, effective, and efficient blood culture Collection Kit from C-Quest Medical. The Kit's intuitive Minimize false positive blood culture results and

culture results.

consistently accurate blood

## As easy as 1, 2, 3

assures reduced false positive readings, and Popular, easy-to-use components require virtually no while the sterile initial specimen diversion tube Everything is in one place, so your time is optimized, training, and customized components are also available lection, Collection, and Post Collection applications. Three pockets separate the Kit's contents into Pre-Col

# COLLECTION POST COLLECTION

# Features and Benefits:

- blood culture contamination. Sterile Initial Specimen Diversion Tube minimizes
- Proper patient diagnosis and treatment are enhanced
- Antibiotic use and length of hospital stay are reduced
- 3-Pocket Kit configuration optimizes your efficiency. without diversion device. Costs no more than alternative collection methods
- Familiar components are easy-to-use and require virtually no training.
- Custom Kit components are available upon request





MINIMUM MINIMUM

Surgical Sponges 8 公公公司

### **BLOOD CULTURE COLLECTION PROCEDURE KIT**

Including Sterile Initial Specimen Diversion Tube

ORDERING INFORMATION		
PART NUMBER	DESCRIPTION	QUANTITY
40-300	Sterile Initial Specimen Diversion Tube	480
40-301	MANUAL - 21g x 3/4" Needle, 12" Butterfly attached to Thin Neck Blood Culture Vial Holder, Sterile (internal & external) Initial Specimen Diversion Tube, Drape, Tourniquet, Hand Sanitizer, Gloves, 2 Alcohol Prep Pads, Chlorhexidine Gluconate, Gauze, Bandage, Biohazard Bag	25
40-302	AUTOMATIC - 21g x 3/4" Needle, 12" Butterfly attached to Thin Neck Blood Culture Vial Holder, Sterile (internal & external) Initial Specimen Diversion Tube, Drape, Tourniquet, Hand Sanitizer, Gloves, 2 Alcohol Prep Pads, Chlorhexidine Gluconate, Gauze, Bandage, Biohazard Bag	25
40-303	MANUAL - 23g x 3/4" Needle, 12" Butterfly attached to Thin Neck Blood Culture Vial Holder, Sterile (internal & external) Initial Specimen Diversion Tube, Drape, Tourniquet, Hand Sanitizer, Gloves, 2 Alcohol Prep Pads, Chlorhexidine Gluconate, Gauze, Bandage, Biohazard Bag	25
40-304	AUTOMATIC - 23g x 3/4" Needle, 12" Butterfly attached to Thin Neck Blood Culture Vial Holder, Sterile (internal & external) Initial Specimen Diversion Tube, Drape, Tourniquet, Hand Sanitizer, Gloves, 2 Alcohol Prep Pads, Chlorhexidine Gluconate, Gauze, Bandage, Biohazard Bag	25
40-401	MANUAL - 21g x 3/4" Needle, 12" Butterfly attached to Wide Neck Blood Culture Vial Holder, Sterile (internal & external) Initial Specimen Diversion Tube, Drape, Tourniquet, Hand Sanitizer, Gloves, 2 Alcohol Prep Pads, Chlorhexidine Gluconate, Gauze, Bandage, Biohazard Bag	25
40-402	AUTOMATIC - 21g x 3/4" Needle, 12" Butterfly attached to Wide Neck Blood Culture Vial Holder, Sterile (internal & external) Initial Specimen Diversion Tube, Drape, Tourniquet, Hand Sanitizer, Gloves, 2 Alcohol Prep Pads, Chlorhexidine Gluconate, Gauze, Bandage, Biohazard Bag	25
40-403	MANUAL - 23g x 3/4" Needle, 12" Butterfly attached to Wide Neck Blood Culture Vial Holder, Sterile (internal & external) Initial Specimen Diversion Tube, Drape, Tourniquet, Hand Sanitizer, Gloves, 2 Alcohol Prep Pads, Chlorhexidine Gluconate, Gauze, Bandage, Biohazard Bag	25
40-404	AUTOMATIC - 23g x 3/4" Needle, 12" Butterfly attached to Wide Neck Blood Culture Vial Holder, Sterile (internal & external) Initial Specimen Diversion Tube, Drape, Tourniquet, Hand Sanitizer, Gloves, 2 Alcohol Prep Pads, Chlorhexidine Gluconate, Gauze, Bandage, Biohazard Bag	25



88 Inverness Circle East, Suite E-102 Englewood, CO 80112 888-617-8217 www.c-questmedical.com



### References

- 1. Rupp ME, Cavalieri RJ, Marolf C, Lyden E. Reduction in Blood Culture Contamination Through Use of Initial Specimen Diversion Device. Clin Infect Dis 2017; 65:201.
- 2. Bell M, Bogar C, Plante J, et al. Effectiveness of a Novel Specimen Collection System in Reducing Blood Culture Contamination Rates. J Emerg Nurs 2018; 44:570.
- 3. Zimmerman FS, Assous MV, Zevin S, Wiener-Well Y. Reducing blood culture contamination using an initial specimen diversion device. Am J Infect Control 2019; 47:822.
- Zimmerman FS, Karameh H, Ben-Chetrit E, et al. Modification of Blood Test Draw Order to Reduce Blood Culture Contamination: A Randomized Clinical Trial. Clin Infect Dis 2020; 71:1215.
- Doern GV, Carroll KC, Diekema DJ, Garey KW, Rupp ME, Weinstein MP, et al. 2020. A Comprehensive Update on the Problem of Blood Culture Contamination and a Discussion of Methods for Addressing the Problem. Clin Microbiol Rev 33: e00009-19.